arXiv:1604.00996v1 [math.FA] 4 Apr 2016

SOME REVERSED AND REFINED CALLEBAUT INEQUALITIES VIA KONTOROVICH CONSTANT

MOJTABA BAKHERAD

Abstract. In this paper we employ some operator techniques to establish some refinements and reverses of the Callebaut inequality involving the geometric mean and Hadamard product under some mild conditions. In particular, we show

$$K\left(\frac{M^{2t-1}}{m^{2t-1}},2\right)^{r'}\sum_{j=1}^{n}(A_j\sharp_s B_j)\circ\sum_{j=1}^{n}(A_j\sharp_{1-s}B_j)$$
$$+\left(\frac{t-s}{t-1/2}\right)\left(\sum_{j=1}^{n}(A_j\sharp_t B_j)\circ\sum_{j=1}^{n}(A_j\sharp_{1-t}B_j)-\sum_{j=1}^{n}(A_j\sharp B_j)\circ\sum_{j=1}^{n}(A_j\sharp B_j)\right)$$
$$\leq\sum_{j=1}^{n}(A_j\sharp_t B_j)\circ\sum_{j=1}^{n}(A_j\sharp_{1-t}B_j),$$

where $A_j,B_j\in\mathbb{B}(\mathscr{H})$ $(1\leq j\leq n)$ are positive operators such that $0<m'\leq B_j\leq m<M\leq A_j\leq M'$ $(1\leq j\leq n)$, either $1\geq t\geq s>\frac{1}{2}$ or $0\leq t\leq s<\frac{1}{2}$, $r'=\min\left\{\frac{t-s}{t-1/2},\frac{s-1/2}{t-1/2}\right\}$ and $K(t,2)=\frac{(t+1)^2}{4t}$ $(t>0)$.

1. Introduction and preliminaries

Let $\mathbb{B}(\mathscr{H})$ denote the C^*-algebra of all bounded linear operators on a complex Hilbert space $\mathscr{H}$ with the identity I. An operator $A\in\mathbb{B}(\mathscr{H})$ is called positive if $\langle Ax,x\rangle\geq 0$ for all $x\in\mathscr{H}$, and we then write $A\geq 0$. We write $A>0$ if A is a positive invertible operator. The set of all positive invertible operators is denoted by $\mathbb{B}(\mathscr{H})_+$. For self-adjoint operators $A,B\in\mathbb{B}(\mathscr{H})$, we say $B\geq A$ if $B-A\geq 0$. The Gelfand map $f\mapsto f(A)$ is an isometric $*$-isomorphism between the C^*-algebra $C(\mathrm{sp(A)})$ of a complex-valued continuous functions on the spectrum sp(A) of a self-adjoint operator A and the C^*-algebra generated by I and A. If $f,g\in C(\mathrm{sp}(A))$, then $f(t)\geq g(t)$ $(t\in\mathrm{sp}(A))$ implies that $f(A)\geq g(A)$.

It is known that the Hadamard product can be presented by filtering the tensor product $A\otimes B$ through a positive linear map. In fact, $A\circ B=U^*(A\otimes B)U$, where

2010 *Mathematics Subject Classification.* Primary 47A63, Secondary 47A60.

Key words and phrases. Callebaut inequality; Cauchy–Schwarz inequality; Hadamard product; Operator geometric mean; Kontorovich constant.

$U : \mathscr{H} \to \mathscr{H} \otimes \mathscr{H}$ is the isometry defined by $Ue_j = e_j \otimes e_j$, where (e_j) is an orthonormal basis of the Hilbert space $\mathscr{H}$; see [7].

For $A, B \in \mathbb{B}(\mathscr{H})_+$, the operator geometric mean $A\sharp B$ is defined by $A\sharp B = A^{\frac{1}{2}}\left(A^{\frac{-1}{2}}BA^{\frac{-1}{2}}\right)^{\frac{1}{2}}A^{\frac{1}{2}}$. For $\alpha \in (0,1)$, the operator weighted geometric mean is defined by

$$A\sharp_\alpha B = A^{\frac{1}{2}}\left(A^{\frac{-1}{2}}BA^{\frac{-1}{2}}\right)^{\alpha}A^{\frac{1}{2}}.$$

Callebaut [4] showed the following refinement of the Cauchy–Schwarz inequality

$$\begin{aligned}\left(\sum_{j=1}^n x_j^{\frac{1}{2}}y_j^{\frac{1}{2}}\right)^2 &\leq \sum_{j=1}^n x_j^{\frac{1+s}{2}}y_j^{\frac{1-s}{2}}\sum_{j=1}^n x_j^{\frac{1-s}{2}}y_j^{\frac{1+s}{2}}\\ &\leq \sum_{j=1}^n x_j^{\frac{1+t}{2}}y_j^{\frac{1-t}{2}}\sum_{j=1}^n x_j^{\frac{1-t}{2}}y_j^{\frac{1+t}{2}}\\ &\leq \left(\sum_{j=1}^n x_j\right)\left(\sum_{j=1}^n y_j\right), \end{aligned} \tag{1.1}$$

where x_j, y_j $(1 \leq j \leq n)$ are positive real numbers and either $1 \geq t \geq s > \frac{1}{2}$ or $0 \leq t \leq s < \frac{1}{2}$. This is indeed an extension of the Cauchy–Schwarz inequality.
Wada [10] gave an operator version of the Callebaut inequality by showing that if $A, B \in \mathbb{B}(\mathscr{H})_+$, then

$$\begin{aligned}(A\sharp B)\otimes(A\sharp B) &\leq \frac{1}{2}\left\{(A\sharp_\alpha B)\otimes(A\sharp_{1-\alpha}B)+(A\sharp_{1-\alpha}B)\otimes(A\sharp_\alpha B)\right\}\\ &\leq \frac{1}{2}\left\{(A\otimes B)+(B\otimes A)\right\},\end{aligned}$$

where $\alpha \in [0,1]$. In [6] the authors showed another operator version of the Callebaut inequality

$$\begin{aligned}\sum_{j=1}^n (A_j\sharp B_j)\circ\sum_{j=1}^n (A_j\sharp B_j) &\leq \sum_{j=1}^n (A_j\sharp_s B_j)\circ\sum_{j=1}^n (A_j\sharp_{1-s} B_j)\\ &\leq \sum_{j=1}^n (A_j\sharp_t B_j)\circ\sum_{j=1}^n (A_j\sharp_{1-t} B_j)\\ &\leq \left(\sum_{j=1}^n A_j\right)\circ\left(\sum_{j=1}^n B_j\right), \end{aligned} \tag{1.2}$$

where $A_j, B_j \in \mathbb{B}(\mathscr{H})_+$ $(1 \leq j \leq n)$ and either $1 \geq t \geq s > \frac{1}{2}$ or $0 \leq t \leq s < \frac{1}{2}$. In [3] the authors presented the following refinement of inequality (1.2) as follows

$$\begin{aligned}
&\sum_{j=1}^{n}(A_j \sharp_s B_j) \circ \sum_{j=1}^{n}(A_j \sharp_{1-s} B_j) \\
&\qquad \leq \sum_{j=1}^{n}(A_j \sharp_s B_j) \circ \sum_{j=1}^{n}(A_j \sharp_{1-s} B_j) \\
&\qquad + \left(\frac{t-s}{s-1/2}\right)\left(\sum_{j=1}^{n}(A_j \sharp_s B_j) \circ \sum_{j=1}^{n}(A_j \sharp_{1-s} B_j) - \sum_{j=1}^{n}(A_j \sharp B_j) \circ \sum_{j=1}^{n}(A_j \sharp B_j)\right) \\
&\qquad \leq \sum_{j=1}^{n}(A_j \sharp_t B_j) \circ \sum_{j=1}^{n}(A_j \sharp_{1-t} B_j)\,, \qquad (1.3)
\end{aligned}$$

in which $A_j, B_j \in \mathbb{B}(\mathscr{H})_+$ $(1 \leq j \leq n)$ and either $1 \geq t \geq s > \frac{1}{2}$ or $0 \leq t \leq s < \frac{1}{2}$. There have been obtained several Cauchy–Schwarz type inequalities for Hilbert space operators and matrices; see [1, 2, 5, 8] and references therein.

In this paper, we present some refinements and reverses of the Callebaut inequality involving the weighted geometric mean and Hadamard product of Hilbert space operators.

2. Further refienements of the Callebaut inequality involving Hadamard product

The Kontorovich constant is

$$K(t,2) = \frac{(t+1)^2}{4t} \qquad (t > 0).$$

The classical Young inequality states that

$$a^{\nu} b^{1-\nu} \leq \nu a + (1-\nu) b,$$

where $a, b \geq 0$ and $\nu \in [0,1]$. Recently, Zuo et. al. [11] showed an improvement of the Young inequality as follows:

$$K\left(\sqrt{\frac{a}{b}}, 2\right)^{r} a^{\nu} b^{1-\nu} \leq \nu a + (1-\nu) b,$$

where $a, b > 0$, $\nu \in [0,1]$, $r = \min\{\nu, 1-\nu\}$. Applying this inequality, J. Wu and J. Zhao [9] showed the following refiniment of the Young inequality

$$K\left(\sqrt{\frac{a}{b}}, 2\right)^{r'} a^{\nu} b^{1-\nu} + r\left(\sqrt{a} - \sqrt{b}\right)^2 \leq \nu a + (1-\nu) b, \qquad (2.1)$$

where $a, b > 0$, $\nu \in [0,1] - \left\{\frac{1}{2}\right\}$, $r = \min\{\nu, 1-\nu\}$ and $r' = \min\{2r, 1-2r\}$. Using (2.1) we get the following lemmas.

Lemma 2.1. *Let $a,b>0$ and $\nu\in[0,1]-\left\{\frac{1}{2}\right\}$. Then*

$$K\left(\sqrt{\frac{a}{b}},2\right)^{r'}\left(a^{\nu}b^{1-\nu}+a^{1-\nu}b^{\nu}\right)+2r\left(\sqrt{a}-\sqrt{b}\right)^{2}\leq a+b, \tag{2.2}$$

where $r=\min\{\nu,1-\nu\}$ and $r'=\min\{2r,1-2r\}$.

Lemma 2.2. *Let $0<m'\leq B\leq m<M\leq A\leq M'$ and either $1\geq t\geq s>\frac{1}{2}$ or $0\leq t\leq s<\frac{1}{2}$. Then*

$$\begin{aligned}K\left(\frac{M^{2t-1}}{m^{2t-1}},2\right)^{r'}&\left(A^{s}\otimes B^{1-s}+A^{1-s}\otimes B^{s}\right)\\&+\left(\frac{t-s}{t-1/2}\right)\left(A^{t}\otimes B^{1-t}+A^{1-t}\otimes B^{t}-2(A^{\frac{1}{2}}\otimes B^{\frac{1}{2}})\right)\\&\leq A^{t}\otimes B^{1-t}+A^{1-t}\otimes B^{t},\end{aligned} \tag{2.3}$$

where $r'=\min\left\{\frac{t-s}{t-1/2},\frac{s-1/2}{t-1/2}\right\}$.

Proof. Let $a>0$. If we replace b by a^{-1} and take $\nu=\frac{1-\mu}{2}$ (2.2), then we get

$$K(a,2)^{r'}\left(a^{\mu}+a^{-\mu}\right)+(1-\mu)\left(a+a^{-1}-2\right)\leq a+a^{-1}, \tag{2.4}$$

in which $\mu\in(0,1]$ and $r'=\min\{1-\mu,\mu\}$. Let us fix positive real numbers α,β such that $\beta<\alpha$. It follows from $0<m'\leq B\leq m<M\leq A\leq M'$ that $I\leq h=\left(\frac{M}{m}\right)^{\alpha}\leq A^{\alpha}\otimes B^{-\alpha}\leq h'=\left(\frac{M'}{m'}\right)^{\alpha}$ and $\mathrm{sp}(\mathrm{A}^{\alpha}\otimes\mathrm{B}^{-\alpha})\subseteq[\mathrm{h},\mathrm{h}']\subseteq(1,+\infty)$. Since the Kontorovich constant $\frac{(a+1)^2}{4a}$ is an increasing function on $(1,+\infty)$, by (2.4) we have

$$K\left(\frac{M^{\alpha}}{m^{\alpha}},2\right)^{r'}\left(a^{\mu}+a^{-\mu}\right)+(1-\mu)\left(a+a^{-1}-2\right)\leq a+a^{-1},$$

where $\mu\in(0,1]$ and $r'=\min\{1-\mu,\mu\}$. Using the functional calculus, if we replace a by the operator $A^{\alpha}\otimes B^{-\alpha}$ and μ by $\frac{\beta}{\alpha}$ we have

$$\begin{aligned}K\left(\frac{M^{\alpha}}{m^{\alpha}},2\right)^{r'}&\left(A^{\beta}\otimes B^{-\beta}+A^{-\beta}\otimes B^{\beta}\right)\\&+\left(1-\frac{\beta}{\alpha}\right)\left(A^{\alpha}\otimes B^{-\alpha}+A^{-\alpha}\otimes B^{\alpha}-2I\right)\\&\leq A^{\alpha}\otimes B^{-\alpha}+A^{-\alpha}\otimes B^{\alpha},\end{aligned} \tag{2.5}$$

where $r'=\min\left\{1-\frac{\beta}{\alpha},\frac{\beta}{\alpha}\right\}$. Multiplying both sides of (2.5) by $A^{\frac{1}{2}}\otimes B^{\frac{1}{2}}$ we reach

$$\begin{aligned}K\left(\frac{M^\alpha}{m^\alpha},2\right)^{r'}&\left(A^{1+\beta}\otimes B^{1-\beta}+A^{1-\beta}\otimes B^{1+\beta}\right)\\&+\left(1-\frac{\beta}{\alpha}\right)\left(A^{1+\alpha}\otimes B^{1-\alpha}+A^{1-\alpha}\otimes B^{1+\alpha}-2(A\otimes B)\right)\\&\leq A^{1+\alpha}\otimes B^{1-\alpha}+A^{1-\alpha}\otimes B^{1+\alpha}.\end{aligned}\tag{2.6}$$

Now, if we replace α,β,A,B by $2t-1,2s-1,A^{\frac{1}{2}},B^{\frac{1}{2}}$ respectively, in (2.6), we obtain

$$\begin{aligned}K\left(\frac{M^{2t-1}}{m^{2t-1}},2\right)^{r'}&\left(A^s\otimes B^{1-s}+A^{1-s}\otimes B^s\right)\\&+\left(\frac{t-s}{t-1/2}\right)\left(A^t\otimes B^{1-t}+A^{1-t}\otimes B^t-2(A^{\frac{1}{2}}\otimes B^{\frac{1}{2}})\right)\\&\leq A^t\otimes B^{1-t}+A^{1-t}\otimes B^t\end{aligned}$$

for either $1\geq t\geq s>\frac{1}{2}$ or $0\leq t\leq s<\frac{1}{2}$ and $r'=\min\left\{\frac{t-s}{t-1/2},\frac{s-1/2}{t-1/2}\right\}$. □

We are ready to prove the first result of this section.

Theorem 2.3. *Let $0<m'\leq B_j\leq m<M\leq A_j\leq M'$ $(1\leq j\leq n)$ and either $1\geq t\geq s>\frac{1}{2}$ or $0\leq t\leq s<\frac{1}{2}$. Then*

$$\begin{aligned}K\left(\frac{M^{2t-1}}{m^{2t-1}},2\right)^{r'}&\sum_{j=1}^n(A_j\sharp_s B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-s}B_j)\\&+\left(\frac{t-s}{t-1/2}\right)\left(\sum_{j=1}^n(A_j\sharp_t B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-t}B_j)-\sum_{j=1}^n(A_j\sharp B_j)\circ\sum_{j=1}^n(A_j\sharp B_j)\right)\\&\leq\sum_{j=1}^n(A_j\sharp_t B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-t}B_j)\,,\end{aligned}\tag{2.7}$$

where $r'=\min\left\{\frac{t-s}{t-1/2},\frac{s-1/2}{t-1/2}\right\}$.

Proof. Put $C_j=A_j^{-\frac{1}{2}}B_jA_j^{-\frac{1}{2}}$ $(1\leq j\leq n)$. By inequality (2.3) we get

$$\begin{aligned}K\left(\frac{M^{2t-1}}{m^{2t-1}},2\right)^{r'}&\left(C_j^s\otimes C_i^{1-s}+C_j^{1-s}\otimes C_i^s\right)\\&+\left(\frac{t-s}{t-1/2}\right)\left(C_j^t\otimes C_i^{1-t}+C_j^{1-t}\otimes C_i^t-2\big(C_j^{\frac{1}{2}}\otimes C_i^{\frac{1}{2}}\big)\right)\\&\leq C_j^t\otimes C_i^{1-t}+C_j^{1-t}\otimes C_i^t\qquad(1\leq i,j\leq n).\end{aligned}\tag{2.8}$$

Multiplying both sides of (2.8) by $A_j^{\frac{1}{2}}\otimes A_i^{\frac{1}{2}}$ we get

$$K\left(\frac{M^{2t-1}}{m^{2t-1}},2\right)^{r'}((A_j\sharp_s B_j)\otimes(A_i\sharp_{1-s}B_i)+(A_j\sharp_{1-s}B_j)\otimes(A_i\sharp_s B_i))$$
$$+\left(\frac{t-s}{t-1/2}\right)\Big((A_j\sharp_t B_j)\otimes(A_i\sharp_{1-t}B_i)+(A_j\sharp_{1-t}B_j)$$
$$\otimes(A_i\sharp_t B_i)-2(A_j\sharp B_j)\otimes(A_i\sharp B_i)\Big)$$
$$\leq(A_j\sharp_t B_j)\otimes(A_i\sharp_{1-t}B_i)+(A_j\sharp_{1-t}B_j)\otimes(A_i\sharp_t B_i) \qquad (2.9)$$

for all $1\leq i,j\leq n$. Therefore

$$K\left(\frac{M^{2t-1}}{m^{2t-1}},2\right)^{r'}\left(\sum_{j=1}^n(A_j\sharp_s B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-s}B_j)\right)$$
$$+\left(\frac{t-s}{t-1/2}\right)\left(\sum_{j=1}^n(A_j\sharp_t B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-t}B_j)-\Big(\sum_{j=1}^n A_j\sharp B_j\Big)\circ\Big(\sum_{j=1}^n A_j\sharp B_j\Big)\right)$$
$$=\frac{1}{2}\sum_{i,j=1}^n\Big[K\left(\frac{M^{2t-1}}{m^{2t-1}},2\right)^{r'}\Big((A_j\sharp_s B_j)\circ(A_i\sharp_{1-s}B_i)+(A_j\sharp_{1-s}B_j)\circ(A_i\sharp_s B_i)\Big)$$
$$+\left(\frac{t-s}{t-1/2}\right)((A_j\sharp_t B_j)\circ(A_i\sharp_{1-t}B_i)+(A_j\sharp_{1-t}B_j)\circ(A_i\sharp_t B_i)-2(A_j\sharp B_j)\circ(A_i\sharp B_i))\Big]$$
$$\leq\frac{1}{2}\sum_{i,j=1}^n((A_j\sharp_t B_j)\circ(A_i\sharp_{1-t}B_i)+(A_j\sharp_{1-t}B_j)\circ(A_i\sharp_t B_i)) \qquad \text{(by inequality (2.9))}$$
$$=\sum_{j=1}^n(A_j\sharp_t B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-t}B_j)\,.$$

□

Remark 2.4. It follows from

$$\left(\frac{t-s}{t-1/2}\right)\left(\sum_{j=1}^n(A_j\sharp_t B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-t}B_j)-\sum_{j=1}^n(A_j\sharp B_j)\circ\sum_{j=1}^n(A_j\sharp B_j)\right)\geq 0\,,$$

where $A_j,B_j\in\mathbb{B}(\mathscr{H})_+$ $(1\leq j\leq n)$, either $1\geq t\geq s>\frac{1}{2}$ or $0\leq t\leq s<\frac{1}{2}$ and $K(t,2)=\frac{(t+1)^2}{4t}\geq 1$ $(t>0)$ that inequality (2.7) is a refinement of the second inequality of inequalities (1.2) and (1.3).

We conclude an application of Theorem 2.3 for numerical cases which is a refinement of inequality (1.1).

Corollary 2.5. *Let $0 < m' \le y_j \le m < M \le x_j \le M'$ $(1 \le j \le n)$ and either $-1 \le t \le s < 0$ or $1 \ge t \ge s > 0$. Then*

$$\begin{aligned}
&\sum_{j=1}^{n} x_j^{\frac{1+s}{2}} y_j^{\frac{1-s}{2}} \sum_{j=1}^{n} x_j^{\frac{1-s}{2}} y_j^{\frac{1+s}{2}} \\
&\le K\left(\frac{M^t}{m^t}, 2\right)^{r'} \sum_{j=1}^{n} x_j^{\frac{1+s}{2}} y_j^{\frac{1-s}{2}} \sum_{j=1}^{n} x_j^{\frac{1-s}{2}} y_j^{\frac{1+s}{2}} \\
&\quad + \left(\frac{t-s}{t}\right)\left(\sum_{j=1}^{n} x_j^{\frac{1+t}{2}} y_j^{\frac{1-t}{2}} \sum_{j=1}^{n} x_j^{\frac{1-t}{2}} y_j^{\frac{1+t}{2}} - \left(\sum_{j=1}^{n} x_j^{\frac{1}{2}} y_j^{\frac{1}{2}}\right)^2\right) \\
&\le \sum_{j=1}^{n} x_j^{\frac{1+t}{2}} y_j^{\frac{1-t}{2}} \sum_{j=1}^{n} x_j^{\frac{1-t}{2}} y_j^{\frac{1+t}{2}},
\end{aligned}$$

where $r' = \min\left\{\frac{t-s}{t}, \frac{s}{t}\right\}$.

Proof. If we put $A_j = x_j, B_j = y_j$ $(1 \le j \le n)$ in Theorem 2.3 and inequality (1.2), then we get

$$\begin{aligned}
&\sum_{j=1}^{n} x_j^{1-s} y_j^{s} \sum_{j=1}^{n} x_j^{s} y_j^{1-s} \\
&\le K\left(\frac{M^{2t-1}}{m^{2t-1}}, 2\right)^{r'} \sum_{j=1}^{n} x_j^{1-s} y_j^{s} \sum_{j=1}^{n} x_j^{s} y_j^{1-s} \\
&\quad + \left(\frac{t-s}{t-1/2}\right)\left(\sum_{j=1}^{n} x_j^{1-t} y_j^{t} \sum_{j=1}^{n} x_j^{t} y_j^{1-t} - \left(\sum_{j=1}^{n} x_j^{\frac{1}{2}} y_j^{\frac{1}{2}}\right)^2\right) \\
&\le \sum_{j=1}^{n} x_j^{1-t} y_j^{t} \sum_{j=1}^{n} x_j^{t} y_j^{1-t},
\end{aligned}$$

where $r' = \min\left\{\frac{t-s}{t-1/2}, \frac{s-1/2}{t-1/2}\right\}$ and either $0 \le t \le s < \frac{1}{2}$ or $1 \ge t \ge s > \frac{1}{2}$. Now if we replace s by $\frac{s+1}{2}$ and t by $\frac{t+1}{2}$, respectively, where either $-1 \le t \le s < 0$ or $1 \ge t \ge s > 0$, then we reach the desired inequalities. □

Lemma 2.6. *Let $a, b > 0$ and $\nu \in (0,1)$. Then*

$$a^{\nu} b^{1-\nu} + a^{1-\nu} b^{\nu} + 2r(\sqrt{a} - \sqrt{b})^2 + r'\left(2\sqrt{ab} + a + b - 2a^{\frac{1}{4}} b^{\frac{3}{4}} - 2a^{\frac{3}{4}} b^{\frac{1}{4}}\right) \le a + b,$$

where $r = \min\{\nu, 1-\nu\}$ and $r' = \min\{2r, 1-2r\}$.

Proof. Let $a,b>0$, $\nu\in(0,1)$, $r=\min\{\nu,1-\nu\}$ and $r'=\min\{2r,1-2r\}$. Applying [12, Lemma 1], we have the inequalities

$$a^{1-\nu}b^{\nu}+\nu(\sqrt{a}-\sqrt{b})^2+r'(\sqrt[4]{ab}-\sqrt{a})^2\leq(1-\nu)a+\nu b\,,$$

where $0<\nu\leq\frac{1}{2}$ and

$$a^{1-\nu}b^{\nu}+(1-\nu)(\sqrt{a}-\sqrt{b})^2+r'(\sqrt[4]{ab}-\sqrt{b})^2\leq(1-\nu)a+\nu b\,,$$

where $\frac{1}{2}<\nu<1$. Summing these inequalities we get the desired result. □

Lemma 2.7. *Let $A_j,B_j\in\mathbb{B}(\mathscr{H})_+$ $(1\leq j\leq n)$ and either $1\geq t\geq s>\frac{1}{2}$ or $0\leq t\leq s<\frac{1}{2}$. Then*

$$\begin{aligned}
&(A^s\otimes B^{1-s}+A^{1-s}\otimes B^s)\\
&\quad+\left(\frac{t-s}{t-1/2}\right)\left(A^s\otimes B^{1-s}+A^{1-s}\otimes B^s-2(A^{\frac{1}{2}}\otimes B^{\frac{1}{2}})\right)\\
&\quad+r'\left(A^t\otimes B^{1-s}+A^{1-s}\otimes B^s+2(A^{\frac{1}{2}}\otimes B^{\frac{1}{2}})-2A^{\frac{1+2s}{4}}\otimes B^{\frac{3-2s}{4}}-2A^{\frac{3-2s}{4}}\otimes B^{\frac{1+2s}{4}}\right)\\
&\leq A^t\otimes B^{1-t}+A^{1-t}\otimes B^t\,,
\end{aligned}$$

where $r'=\min\left\{\frac{t-s}{t-1/2},\frac{s-1/2}{t-1/2}\right\}$.

Using Lemma 2.7 and the same argument in the proof of Lemma 2.2, we get another refinement of inequality (1.2).

Theorem 2.8. *Let $A_j,B_j\in\mathbb{B}(\mathscr{H})_+$ $(1\leq j\leq n)$ and either $1\geq t\geq s>\frac{1}{2}$ or $0\leq t\leq s<\frac{1}{2}$. Then*

$$\begin{aligned}
&\sum_{j=1}^n(A_j\sharp_s B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-s}B_j)\\
&\quad+\left(\frac{t-s}{t-1/2}\right)\left(\sum_{j=1}^n(A_j\sharp_s B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-s}B_j)-\sum_{j=1}^n(A_j\sharp B_j)\circ\sum_{j=1}^n(A_j\sharp B_j)\right)\\
&\quad+r'\Big(\sum_{j=1}^n(A_j\sharp_s B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-s}B_j)+\sum_{j=1}^n(A_j\sharp B_j)\circ\sum_{j=1}^n(A_j\sharp B_j)\\
&\quad-2\sum_{j=1}^n(A_j\sharp_{\frac{3-2s}{4}}B_j)\circ\sum_{j=1}^n(A_j\sharp_{\frac{1+2s}{4}}B_j)\Big)\\
&\leq\sum_{j=1}^n(A_j\sharp_t B_j)\circ\sum_{j=1}^n(A_j\sharp_{1-t}B_j)\,,
\end{aligned}\tag{2.10}$$

where $r'=\min\left\{\frac{t-s}{t-1/2},\frac{s-1/2}{t-1/2}\right\}$.

Remark 2.9. If $A_j, B_j \in \mathbb{B}(\mathscr{H})_+$ $(1 \le j \le n)$ and either $1 \ge t \ge s > \frac{1}{2}$ or $0 \le t \le s < \frac{1}{2}$, then

$$\sum_{j=1}^{n}(A_j \sharp_s B_j) \circ \sum_{j=1}^{n}(A_j \sharp_{1-s} B_j) + \sum_{j=1}^{n}(A_j \sharp B_j) \circ \sum_{j=1}^{n}(A_j \sharp B_j) - 2\sum_{j=1}^{n}(A_j \sharp_{\frac{3-2s}{4}} B_j) \circ \sum_{j=1}^{n}(A_j \sharp_{\frac{1+2s}{4}} B_j)$$

and

$$\left(\frac{t-s}{t-1/2}\right)\left(\sum_{j=1}^{n}(A_j \sharp_s B_j) \circ \sum_{j=1}^{n}(A_j \sharp_{1-s} B_j) - \sum_{j=1}^{n}(A_j \sharp B_j) \circ \sum_{j=1}^{n}(A_j \sharp B_j)\right)$$

are positive operators. So inequality (2.10) is another refinement of the second inequality of inequalities (1.2) and (1.3).

If we put $B_j = I$ $(1 \le j \le n)$ in Theorem 2.8, then we get the next result.

Corollary 2.10. *Let $A_j \in \mathbb{B}(\mathscr{H})_+$ $(1 \le j \le n)$ and either $1 \ge t \ge s > \frac{1}{2}$ or $0 \le t \le s < \frac{1}{2}$. Then*

$$\begin{aligned}
&\sum_{j=1}^{n} A_j^{1-s} \circ \sum_{j=1}^{n} A_j^{s} \\
&+ \left(\frac{t-s}{t-1/2}\right)\left(\sum_{j=1}^{n} A_j^{1-s} \circ \sum_{j=1}^{n} A_j^{s} - \sum_{j=1}^{n} A_j^{\frac{1}{2}} \circ \sum_{j=1}^{n} A_j^{\frac{1}{2}}\right) \\
&+ r'\Big(\sum_{j=1}^{n} A_j^{1-s} \circ \sum_{j=1}^{n} A_j^{s} + \sum_{j=1}^{n} A_j^{\frac{1}{2}} \circ \sum_{j=1}^{n} A_j^{\frac{1}{2}} \\
&- 2\sum_{j=1}^{n} A_j^{\frac{1+2s}{4}} \circ \sum_{j=1}^{n} A_j^{\frac{3-2s}{4}}\Big) \\
&\le \sum_{j=1}^{n} A_j^{1-t} \circ \sum_{j=1}^{n} A_j^{t},
\end{aligned}$$

where $r' = \min\left\{\frac{t-s}{t-1/2}, \frac{s-1/2}{t-1/2}\right\}$.

If in Theorem 2.8 we replace $A_j, B_j, s, t,$ by $x_j, y_j, \frac{s+1}{2}, \frac{t+1}{2}$ $(1 \le j \le n)$, respectively, then we reach another refinement of inequality (1.1).

Corollary 2.11. *Let x_j, y_j $(1 \leq j \leq n)$ be positive numbers and either $-1 \leq t \leq s < 0$ or $1 \geq t \geq s > 0$. Then*

$$\begin{aligned}
&\sum_{j=1}^{n} x_j^{\frac{1+s}{2}} y_j^{\frac{1-s}{2}} \sum_{j=1}^{n} x_j^{\frac{1-s}{2}} y_j^{\frac{1+s}{2}} \\
&+ \left(\frac{t-s}{t}\right)\left(\sum_{j=1}^{n} x_j^{\frac{1+s}{2}} y_j^{\frac{1-s}{2}} \sum_{j=1}^{n} x_j^{\frac{1-s}{2}} y_j^{\frac{1+s}{2}} - \left(\sum_{j=1}^{n} x_j^{\frac{1}{2}} y_j^{\frac{1}{2}}\right)^2\right) \\
&+ r'\Big(\sum_{j=1}^{n} x_j^{\frac{1+s}{2}} y_j^{\frac{1-s}{2}} \sum_{j=1}^{n} x_j^{\frac{1-s}{2}} y_j^{\frac{1+s}{2}} + \left(\sum_{j=1}^{n} x_j^{\frac{1}{2}} y_j^{\frac{1}{2}}\right)^2 \\
&- 2\sum_{j=1}^{n} x_j^{\frac{2+s}{4}} y_j^{\frac{2-s}{4}} \sum_{j=1}^{n} x_j^{\frac{2-s}{4}} y_j^{\frac{2+s}{4}}\Big) \\
&\leq \sum_{j=1}^{n} x_j^{\frac{1+t}{2}} y_j^{\frac{1-t}{2}} \sum_{j=1}^{n} x_j^{\frac{1-t}{2}} y_j^{\frac{1+t}{2}},
\end{aligned}$$

where $r' = \min\left\{\frac{t-s}{t}, \frac{s-1}{t-1}\right\}$.

3. Some reverses of the Callebaut type inequality

In [9], the authors showed a reverse of the young inequality as follows:

$$\nu a + (1-\nu)b \leq K\left(\sqrt{\frac{a}{b}}, 2\right)^{-r'} a^{\nu} b^{1-\nu} + s\left(\sqrt{a} - \sqrt{b}\right)^2, \tag{3.1}$$

in which $a, b > 0$, $\nu \in [0,1] - \left\{\frac{1}{2}\right\}$, $r = \min\{\nu, 1-\nu\}$, $r' = \min\{2r, 1-2r\}$ and $s = \max\{\nu, 1-\nu\}$. Applying (3.1) we have the next result.

Lemma 3.1. *Let $a, b > 0$ and $\nu \in [0,1] - \left\{\frac{1}{2}\right\}$. Then*

$$a + b \leq K\left(\sqrt{\frac{a}{b}}, 2\right)^{-r'} \left(a^{\nu} b^{1-\nu} + a^{1-\nu} b^{\nu}\right) + 2s\left(\sqrt{a} - \sqrt{b}\right)^2,$$

where $r = \min\{\nu, 1-\nu\}$, $r' = \min\{2r, 1-2r\}$ and $s = \max\{\nu, 1-\nu\}$.
In particular, If $\nu \in [0, \frac{1}{2})$, then

$$a + a^{-1} \leq K(a,2)^{-r'}\left(a^{1-2\nu} + a^{-(1-2\nu)}\right) + 2(1-\nu)\left(a^{\frac{1}{2}} - a^{\frac{-1}{2}}\right)^2. \tag{3.2}$$

Now, utilizing inequality (3.2) and the same argument in the proof of Lemma 2.2 we can accomplish the corresponding result.

Lemma 3.2. *Let $0 < m' \le B \le m < M \le A \le M'$ and either $1 \ge t \ge s > \frac{1}{2}$ or $0 \le t \le s < \frac{1}{2}$. Then*

$$A^t \otimes B^{1-t} + A^{1-t} \otimes B^t \le K\left(\frac{M^{2t-1}}{m^{2t-1}}, 2\right)^{-r'} (A^s \otimes B^{1-s} + A^{1-s} \otimes B^s) + \left(\frac{s-1/2}{t-1/2}\right)\left(A^t \otimes B^{1-t} + A^{1-t} \otimes B^t - 2(A^{\frac{1}{2}} \otimes B^{\frac{1}{2}})\right),$$

where $r' = \min\left\{\frac{t-s}{t-1/2}, \frac{s-1/2}{t-1/2}\right\}$.

As a consequence of Lemma 3.2 we have the following result.

Theorem 3.3. *Let $0 < m' \le B_j \le m < M \le A_j \le M'$ $(1 \le j \le n)$. Then*

$$\sum_{j=1}^n (A_j \sharp_t B_j) \circ \sum_{j=1}^n (A_j \sharp_{1-t} B_j) \le K\left(\frac{M^{2t-1}}{m^{2t-1}}, 2\right)^{-r'} \sum_{j=1}^n (A_j \sharp_s B_j) \circ \sum_{j=1}^n (A_j \sharp_{1-s} B_j) + \left(\frac{s-1/2}{t-1/2}\right)\left(\sum_{j=1}^n (A_j \sharp_t B_j) \circ \sum_{j=1}^n (A_j \sharp_{1-t} B_j) - \sum_{j=1}^n (A_j \sharp B_j) \circ \sum_{j=1}^n (A_j \sharp B_j)\right)$$

for either $1 \ge t \ge s > \frac{1}{2}$ or $0 \le t \le s < \frac{1}{2}$ and $r' = \min\left\{\frac{t-s}{t-1/2}, \frac{s-1/2}{t-1/2}\right\}$.

Remark 3.4. If we put $t = 1$ and $1 \ge s > \frac{1}{2}$ in Theorem 3.3, then we get a reverse of the third inequality of (1.2)

$$\left(\sum_{j=1}^n A_j\right) \circ \left(\sum_{j=1}^n B_j\right) \le K\left(\frac{M}{m}, 2\right)^{-r'} \sum_{j=1}^n (A_j \sharp_s B_j) \circ \sum_{j=1}^n (A_j \sharp_{1-s} B_j) + (2s-1)\left(\left(\sum_{j=1}^n A_j\right) \circ \left(\sum_{j=1}^n B_j\right) - \sum_{j=1}^n (A_j \sharp B_j) \circ \sum_{j=1}^n (A_j \sharp B_j)\right),$$

where $r' = \min\{2-2s, 2s-1\}$.

If we replace $A_j, B_j, s, t,$ by $x_j, y_j, \frac{s+1}{2}, \frac{t+1}{2}$ $(1 \le j \le n)$ in Theorem 3.3, respectively, then we reach a reverse of the second inequality of (1.1)

Corollary 3.5. *Let $0 < m' \le y_j \le m < M \le x_j \le M'$ $(1 \le j \le n)$. Then*

$$\sum_{j=1}^n x_j^{\frac{1+t}{2}} y_j^{\frac{1-t}{2}} \sum_{j=1}^n x_j^{\frac{1-t}{2}} y_j^{\frac{1+t}{2}} \le K\left(\frac{M^{2t-1}}{m^{2t-1}}, 2\right)^{-r'} \sum_{j=1}^n x_j^{\frac{1+s}{2}} y_j^{\frac{1-s}{2}} \sum_{j=1}^n x_j^{\frac{1-s}{2}} y_j^{\frac{1+s}{2}} + \left(\frac{s-1/2}{t-1/2}\right)\left(\sum_{j=1}^n x_j^{\frac{1+t}{2}} y_j^{\frac{1-t}{2}} \sum_{j=1}^n x_j^{\frac{1-t}{2}} y_j^{\frac{1+t}{2}} - \left(\sum_{j=1}^n x_j^{\frac{1}{2}} y_j^{\frac{1}{2}}\right)^2\right)$$

for either $1 \ge t \ge s > \frac{1}{2}$ or $0 \le t \le s < \frac{1}{2}$ and $r' = \min\left\{\frac{t-s}{t-1/2}, \frac{s-1/2}{t-1/2}\right\}$.

Proposition 3.6. *Let $0 < m' \leq B_j \leq m < M \leq A_j \leq M'$ $(1 \leq j \leq n)$ and either $1 \geq t \geq s > \frac{1}{2}$ or $0 \leq t \leq s < \frac{1}{2}$. Then*

$$\begin{aligned} K\left(h^{2t-1},2\right)^{r'} \sum_{j=1}^{n}(A_j\sharp_s B_j)\circ\sum_{j=1}^{n}(A_j\sharp_{1-s}B_j) &+ \left(\frac{t-s}{t-1/2}\right)\left(\sqrt{h}-\sqrt{\frac{1}{h}}\right)^2 \\ &\leq \sum_{j=1}^{n}(A_j\sharp_t B_j)\circ\sum_{j=1}^{n}(A_j\sharp_{1-t}B_j) \\ &\leq K\left(h^{2t-1},2\right)^{-r'}\sum_{j=1}^{n}(A_j\sharp_s B_j)\circ\sum_{j=1}^{n}(A_j\sharp_{1-s}B_j) \\ &\quad + \left(\frac{s-1/2}{t-1/2}\right)\left(\sqrt{h'}-\sqrt{\frac{1}{h'}}\right)^2, \end{aligned}$$

where $h = \frac{M}{m}$, $h' = \frac{M'}{m'}$ and $r' = \min\left\{\frac{t-s}{t-1/2}, \frac{s-1/2}{t-1/2}\right\}$.

Proof. Since the function $f(a) = a - \frac{1}{a}$ is increasing on $(0,\infty)$, we have

$$\left(\sqrt{h}-\sqrt{\frac{1}{h}}\right)^2 \leq \left(\sqrt{a}-\sqrt{\frac{1}{a}}\right)^2 \leq \left(\sqrt{h'}-\sqrt{\frac{1}{h'}}\right)^2 \quad (h \leq a \leq h').$$

Applying inequalities (2.2), (3.2) and the same argument in the proof of Theorem 2.3 we get the desired result. □

Acknowledgement. The author would like to sincerely thank the anonymous referee for some useful comments and suggestions. The author also would like to thank the Tusi Mathematical Research Group (TMRG).

References

[1] J.M. Aldaz, S. Barza, M. Fujii and M.S. Moslehian, *Advances in operator Cauchy–Schwarz inequalities and their reverses*. Ann. Funct. Anal. **6** (2015), no. 3, 275–295.

[2] Lj. Arambasić, D. Bakić and M.S. Moslehian, *A treatment of the Cauchy–Schwarz inequality in C^*-modules*, J. Math. Anal. Appl. **381** (2011) 546–556.

[3] M. Bakherad and M.S. Moslehian, *Complementary and refined inequalities of Callebaut inequality*, Linear Multilinear Algebra **63** (2015), no. 8, 1678–1692.

[4] D.K. Callebaut, *Generalization of the Cauchy–Schwarz inequality*, J. Math. Anal. Appl. **12** (1965), 491-494.

[5] D. Ilišević and S. Varošanec, *On the Cauchy–Schwarz inequality and its reverse in semi-inner product C^*-modules*, Banach J. Math. Anal. **1** (2007), 78–84.

[6] M.S. Moslehian, J.S. Matharu and J.S. Aujla, *Non-commutative Callebaut inequality*, Linear Algebra Appl. **436** (2012) no. 9, 3347–3353.

[7] V.I. Paulsen, *Completely bounded maps and dilation*, Pitman Research Notes in Mathematics Series, **146**, John Wiley & Sons, Inc., New York, 1986.

[8] A. Salemi and A. Sheikh Hosseini, *On reversing of the modified Young inequality*, Ann. Funct. Anal. **5** (2014), no. 1, 70–76.

[9] J. Wu and J. Zhao, *Operator inequalities and reverse inequalities related to the Kittaneh–Manasrah inequalities*, Linear Multilinear Algebra **62** (2014), no. 7, 884–894.

[10] S. Wada, *On some refinement of the Cauchy–Schwarz inequality*, Linear Algebra Appl. **420** (2007) no. 2-3, 433–440.

[11] H. Zuo, G. Shi and M. Fujii, *Refined Young inequality with Kantorovich constant*, J. Math. Inequal., 2011, **5**, 551–556.

[12] J. Zhao and J. Wu, *Operator inequalities involving improved Young and its reverse inequalities*, J. Math. Anal. Appl. **421** (2015), no. 2, 1779–1789.

Department of Mathematics, Faculty of Mathematics, University of Sistan and Baluchestan, Zahedan, Iran.

E-mail address: mojtaba.bakherad@yahoo.com; bakherad@member.ams.org